

Math 211
Final Solutions
December 2001

1) Consider the ordinary differential equation

$$yy' = 2t\sqrt{y^2 + 4}.$$

(a)(4p) Find the general solution.

Separate the variables and integrate:

$$\int \frac{y}{\sqrt{y^2 + 4}} dy = \int 2t dt,$$

so

$$\sqrt{y^2 + 4} = t^2 + C$$

and we obtain the general solution

$$\boxed{y^2 = (t^2 + C)^2 - 4.}$$

(b)(4p) Find the solution of the initial value problem when $y(2) = -\sqrt{5}$.

Using the previous part we obtain $C = -1$ and then

$$\boxed{y = -\sqrt{(t^2 - 1)^2 - 4.}$$

(c)(4p) Find the interval of existence of this solution.

We need $(t^2 - 1)^2 > 4$, i.e. $t^2 - 1 > 2$ or $t^2 > 3$ and the interval contains 2, so it is $\boxed{[\sqrt{3}, \infty)}$.

2)(10p) Find the general solution of the ordinary differential equation.

$$t^2 y' + 3ty = \frac{\sin t}{t} \quad (t > 0).$$

The integrating factor is

$$e^{\int \frac{3}{t} dt} = e^{3 \ln t} = e^{\ln t^3} = t^3,$$

the differential equation becomes

$$t^3 y' + 3t^2 y = \sin t,$$

so

$$(t^3 y)' = \sin t$$

and then

$$t^3 y = -\cos t + C$$

and we obtain the general solution

$$\boxed{y = \frac{C - \cos t}{t^3}.}$$

3)(10p) Three populations exist together and interact in isolated circumstances. Denote the populations by $x_1(t)$, $x_2(t)$ and $x_3(t)$. They interact as follows:

- The first population preys upon the other two and would decline dramatically if $x_2 = x_3 = 0$.
- The second population preys upon the third and would be able to survive if the third were not present, but its growth would be limited by the resources.
- The third population would flourish in the absence of the other two, but its growth would be limited by the resources.

Model the interactions between the three populations with a system of ordinary differential equations. You are **not** required to solve the equations.

One possible model is

$$\begin{aligned}x_1' &= (-a + bx_2 + cx_3)x_1 \\x_2' &= (d - ex_2 - fx_1 + gx_3)x_2 \\x_3' &= (h - ix_3 - jx_1 - kx_2)x_3,\end{aligned}$$

where the constants $a - k$ are all positive.

4) Consider the system of ordinary differential equations

$$y' = Ay, \text{ where } A = \begin{bmatrix} -1 & 1 & 1 \\ -4 & 3 & 2 \\ 0 & 0 & 1 \end{bmatrix}.$$

(a)(3p) Find a fundamental set of solutions.

First we find the eigenvalues and the corresponding eigenvectors. Evaluate the following determinant by the last row:

$$\begin{aligned}\det \begin{bmatrix} -1 - \lambda & 1 & 1 \\ -4 & 3 - \lambda & 2 \\ 0 & 0 & 1 - \lambda \end{bmatrix} &= (1 - \lambda)((-1 - \lambda)(3 - \lambda) + 4) = \\ &= (1 - \lambda)(\lambda^2 - 2\lambda + 1) = (1 - \lambda)(\lambda - 1)^2.\end{aligned}$$

So $\lambda = 1$ is a triple eigenvalue. Eigenvectors:

$$A - I = \begin{bmatrix} -2 & 1 & 1 \\ -4 & 2 & 2 \\ 0 & 0 & 0 \end{bmatrix},$$

and we see that the geometric multiplicity is 2, having two free variables.

Two eigenvectors: $\begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}$. So we have two solutions in the form

$$\boxed{\Phi_1(t) = e^t \begin{bmatrix} 1 \\ 2 \\ 0 \end{bmatrix} \quad \Phi_2(t) = e^t \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix} .}$$

We can find a third by looking at the nullspace of $(A - I)^2 = 0$, so we obtain

that a generalized eigenvector is $v = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$ and a solution is

$$\boxed{\Phi_3(t) = e^t [I + t(A - I)]v = e^t \begin{bmatrix} 1 - 2t \\ -4t \\ 0 \end{bmatrix} .}$$

(b)(3p) Find the general solution of the system.

The general solution is

$$\boxed{y(t) = c_1 \Phi_1(t) + c_2 \Phi_2(t) + c_3 \Phi_3(t) .}$$

(c)(3p) Find the solution of the initial value problem when $y(0) = \begin{bmatrix} 3 \\ 2 \\ 2 \end{bmatrix}$.

We find from the previous part that $y(0) = \begin{bmatrix} c_1 + c_2 + c_3 \\ 2c_1 \\ 2c_2 \end{bmatrix} = \begin{bmatrix} 3 \\ 2 \\ 2 \end{bmatrix}$, so

$c_1 = c_2 = c_3 = 1$ and the solution we look for is

$$\boxed{y(t) = \begin{bmatrix} 3e^t - 2te^t \\ 2e^t - 4te^t \\ 2e^t \end{bmatrix} .}$$

(d)(2p) What can we say about the stability of the equilibrium point at the origin?

The real parts of the eigenvalues are positive, so the equilibrium point is

unstable.

2nd Solution. (a) (b) (c) We have one eigenvalue with multiplicity 3, so we can actually evaluate e^{At} easily. It is easy to check that $(A - I)^2 = 0$, so

$$e^{At} = e^t [I + (A - I)t] = e^t \begin{bmatrix} 1 - 2t & t & t \\ -4t & 1 + 2t & 2t \\ 0 & 0 & 1 \end{bmatrix} .$$

Then the answer to (a): the columns of this matrix. The answer to (b): $e^{At}c$

where $c \in \mathbb{R}^3$. The answer to (c): $e^{At} \begin{bmatrix} 3 \\ 2 \\ 2 \end{bmatrix}$.

5) Consider the periodically forced harmonic ordinary differential equation

$$y'' + 2y' + 2y = 4 \cos 3t.$$

(a)(3p) Find a fundamental set of solutions of the associated homogeneous equation.

Characteristic equation: $\lambda^2 + 2\lambda + 2 = 0$, roots are $\lambda_1 = -1 + i$ and $\lambda_2 = \overline{\lambda_1} = -1 - i$. This means that a fundamental solution set is

$$\Phi_1(t) = e^{-t} \cos t \quad \Phi_2(t) = e^{-t} \sin t.$$

(b)(3p) What is the steady-state solution?

Complex method: the inhomogeneous term is the real part of $4e^{3it}$, so we look for a solution in the form $z(t) = Ae^{3it}$ and the real part of that will be our solution. By plugging $z(t)$ into the equation we obtain

$$(-7 + 6i)Ae^{3it} = 4e^{3it},$$

so $A = \frac{4}{-7+6i} = \frac{4(-7-6i)}{85} = -\frac{28}{85} - \frac{24}{85}i$. This gives us the solution

$$y(t) = -\frac{28}{85} \cos 3t + \frac{24}{85} \sin 3t.$$

Real method: try to find a solution in the form $A \cos 3t + B \sin 3t$. After differentiation we obtain

$$-9A \cos 3t - 9B \sin 3t + 2(-3A \sin 3t + 3B \cos 3t) + 2(A \cos 3t + B \sin 3t) = 4 \cos 3t,$$

so $-7A + 6B = 4$ and $-6A - 7B = 0$. Solving this we get the previous result: $A = -\frac{28}{85}$, $B = \frac{24}{85}$.

(c)(3p) Find the general solution of the inhomogeneous equation.

$$\text{It is } y(t) = -\frac{28}{85} \cos 3t + \frac{24}{85} \sin 3t + c_1 e^{-t} \cos t + c_2 e^{-t} \sin t.$$

(d)(3p) Find the amplitude of the steady-state solution.

It is

$$A = \sqrt{\left(\frac{28}{85}\right)^2 + \left(\frac{24}{85}\right)^2} = \frac{\sqrt{1360}}{85}.$$

6) Consider the system of ordinary differential equations

$$\begin{aligned} x' &= x - xy \\ y' &= y - x^2 + 3. \end{aligned}$$

(a)(3p) Find the nullclines and sketch them, clearly indicating which is the x -nullcline and which is the y -nullcline.

x -nullclines: $x = 0$ (the y -axis) and $y = 1$ (a horizontal line).

y -nullclines: $y = x^2 - 3$ (a parabola).

(b)(3p) Find all of the equilibrium points.

We have three eq. points.

$$\begin{array}{l} x = 0 \\ y = -3 \end{array}$$

$$\begin{array}{l} x = 2 \\ y = 1 \end{array}$$

$$\begin{array}{l} x = -2 \\ y = 1 \end{array}$$

(c)(3p) Compute the Jacobian matrix.

$$J = \begin{bmatrix} 1 - y & -x \\ -2x & 1 \end{bmatrix}.$$

(d)(3p) Classify the equilibrium points.

At $x = 0, y = -3$ the Jacobian is $J = \begin{bmatrix} 4 & 0 \\ 0 & 1 \end{bmatrix}$ and we conclude it is a nodal source.

At $x = 2, y = 0$ the Jacobian is $J = \begin{bmatrix} 1 & -2 \\ -4 & 1 \end{bmatrix}$ and we conclude it is a saddle.

At $x = -2, y = 0$ the Jacobian is $J = \begin{bmatrix} 1 & 2 \\ 4 & 1 \end{bmatrix}$ and we conclude it is a saddle.

7) Consider the system of ordinary differential equations

$$\begin{aligned} x' &= -x \\ y' &= 2y + x^2. \end{aligned}$$

(a)(3p) Show that the y -axis is invariant.

If we start a solution on the y -axis, then the x -coordinate is zero, $x' = -x = 0$, so the x -coordinate will not change, we will stay on the y -axis.

(b)(3p) Show that for any $\begin{bmatrix} c_1 \\ c_2 \end{bmatrix}$

$$\begin{bmatrix} x(t) \\ y(t) \end{bmatrix} = \begin{bmatrix} c_1 e^{-t} \\ c_2 e^{2t} + \frac{c_1^2}{4} (e^{2t} - e^{-2t}) \end{bmatrix}$$

is the solution of the initial value problem with $\begin{bmatrix} x(0) \\ y(0) \end{bmatrix} = \begin{bmatrix} c_1 \\ c_2 \end{bmatrix}$.

Clearly $\begin{bmatrix} x(0) \\ y(0) \end{bmatrix} = \begin{bmatrix} c_1 \\ c_2 \end{bmatrix}$. Also, $x' = -c_1 e^{-t} = -x$ and $y' = 2c_2 e^{2t} + \frac{c_2^2}{2} e^{2t} + \frac{c_1^2}{2} e^{-2t} = 2y + x^2$.

(c)(4p) Show that the set $S = \{(x, y) : y < -\frac{x^2}{2}\}$ is (positively) invariant.

This is the region under the parabola $y = -\frac{x^2}{2}$. On this parabola, $y' = 0$ by the differential equation and $x' > 0$ when $x < 0$ and $x' < 0$ when $x > 0$, so we cannot leave this region. (Also, $x = y = 0$ is an equilibrium point.)

8) Consider the system of ordinary differential equations

$$\begin{aligned} x' &= x(r^3 - 4r^2 + 4) - y \\ y' &= y(r^3 - 4r^2 + 4) + x, \end{aligned}$$

where $r^2 = x^2 + y^2$ (or $r = \sqrt{x^2 + y^2}$).

(a)(5p) Show that the origin is a spiral source.

Change to polar coordinates. The equations become

$$\begin{aligned} r' &= r(r^3 - 4r^2 + 4) \\ \theta' &= 1, \end{aligned}$$

This shows that the origin is a spiral source.

Other possibility: linearize the system, the Jacobian is $\begin{bmatrix} 4 & -1 \\ 1 & 4 \end{bmatrix}$ which shows it is a spiral source again.

(b)(6p) Show that there is a closed solution curve in the annular region

$$A = \{(x, y) : 1 < r < 2\}.$$

From the previous part we can see that this region does not contain an equilibrium point ($\theta' = 1$). Also, the region is positively invariant because $r' = -8 < 0$ when $r = 2$ and $r' = 1 > 0$ when $r = 1$. The assumptions of the Poincaré-Bendixson theorem stand, so the region contains a closed solution curve. (Actually, if we want to be very precise, the assumptions of P-B do not stand, this region is not closed. But we can choose $1 < r_1 < r_2 < 2$ in a way that $r' > 0$ when $r = r_1$ and $r' < 0$ when $r = r_2$ and then apply P-B for $A_1 = \{(x, y) : r_1 \leq r \leq r_2\} \subset A$.)

2nd Solution. For (b): from the polar coordinate form it is clear that there is a value r^* between 1 and 2 such that $r' = 0$ if $r(t) = r^*$. This, together with the equation for θ' proves the statement.

9) Consider the matrix

$$A = \begin{bmatrix} 2 & 7 & 2 \\ 1 & 2 & 0 \\ 0 & 3 & 2 \end{bmatrix}.$$

(a)(4p) Find the nullspace of A .

One possible echelon form of the matrix:

$$A \rightsquigarrow \begin{bmatrix} 1 & 2 & 0 \\ 0 & 3 & 2 \\ 0 & 0 & 0 \end{bmatrix}.$$

We have one free variable, and we obtain that the nullspace is

$$\text{null}(A) = \text{span}\left\{ \begin{bmatrix} 4 \\ -2 \\ 3 \end{bmatrix} \right\}.$$

(b)(4p) What is $\det(A)$?

According to (a), A is singular, so its determinant is 0.

(c)(4p) Are the column vectors in A linearly independent?

According to (a), A is singular, so they are not linearly independent.